



This is a section from the

2022/2023

Mid-Atlantic Commercial Vegetable Production Recommendations

The recommendations are **NOT** for home gardener use.

The **full manual**, containing recommendations specific to New Jersey, can be found on the Rutgers NJAES website in the Publications section: <http://njaes.rutgers.edu/pubs/publication.asp?pid=E001>.

This manual will be revised biennially. **In January 2023, a Critical Update** with important updates to the 2022/2023 manual will be communicated through local Extension Agents and Vegetable Specialists.

The **label** is a legally-binding contract between the user and the manufacturer. The user must follow all rates and restrictions as per label directions. The use of any pesticide inconsistent with the label directions is a violation of federal law.

Cooperating Agencies: Rutgers, The State University of New Jersey, U.S. Department of Agriculture, and County Boards of Commissioners. Rutgers Cooperative Extension, a unit of the Rutgers New Jersey Agricultural Experiment Station, is an equal opportunity program provider and employer.

F. Commodity Recommendations

Pesticide Use Disclaimer

THE LABEL IS THE LAW

Before using a pesticide, check the labeling distributed with the product at the point of sale for legally enforceable rates and use restrictions and precautions. Although labels are available on the Internet from electronic label services such as CDMS (<http://www.cdms.net/>), Greenbook (<https://www.greenbook.net/>), or Agrian (<https://www.agrian.com/labelcenter/results.cfm>) the information contained in these electronic labels may not be identical to the labeling distributed with the product. **Please be advised that these electronic label services provide use disclaimers, and in some cases legally binding *User Agreements* assigning all liability to user of service.** (See section D 3.1. Labels and Labeling for more detail.)

Guide to the Recommended Pesticide Tables in the Following Crop Sections:

1. Pesticides are listed by **group number or code based on chemical structure and mechanism of action**, as classified by the Herbicide Resistance Action Committee (**HRAC**, <https://hracglobal.com/>) for herbicides, the Insecticide Resistance Action Committee (**IRAC**, <https://irac-online.org/>) for insecticides, and the Fungicide Resistance Action Committee (**FRAC**, <https://www.frac.info/>) for fungicides.
In this guide, if the group number or code is in bold font, there are resistance concerns for the product.
2. **Restricted use pesticides** are marked with a * in the Tables. These products may only be used by certified and/or licensed pesticide applicators, and when stated on the label, those making applications under their direct supervision. Some labels may restrict use solely to certified and/or licensed applicators. (See section D 3.2.1 Restricted Use Classification Statement for more detail).
3. **In addition to the pesticide products listed in the Commodity Recommendations below, other formulations or brands with the same active ingredient(s) may be commercially available. ALWAYS CHECK THE INDIVIDUAL PRODUCT LABELING:**
 - a) to ensure a pesticide is labeled for the same intended use,
 - b) to ensure the pesticide is labeled for the desired crop,
 - c) for differences in application rates and % active ingredient(s), and
 - d) additional restrictions.
4. All pesticide recommendations contained in this document are prescribed for spray applications to a **broadcast area of 1 acre** (43,560 square feet). **Adjust the rate accordingly for banded applications** (See section E 1.3. Calibrating Granular Applicators) **or for chemigation** (check labels for amounts per 1,000 feet).
5. Check the label for and do not exceed the maximum amount of pesticide per application and the maximum number of applications per year.
6. **Bee Toxicity Rating (Bee TR):** N=nontoxic; L=minimum impact on bees; M=moderately toxic, can be used if dosage, timing, and method of application are correct, but should NOT be applied directly to the crop if bees are present; H=highly toxic, severe losses expected, -- = data not available.
7. In accordance with the USDA National Organic Program, the Organic Materials Research Institute (OMRI) maintains a directory of all products that OMRI has determined are allowed for use in organic production, processing, and handling. These products are catalogued online in the **OMRI Products List** (see <https://www.omri.org/omri-lists>).

Pumpkins and Winter Squash

Recommended Varieties

Varieties are listed by maturity within each type, earliest first (*=hybrid varieties).

Disease resistance or tolerance in parentheses:

BRT=Black Rot tolerant,

FR=*Fusarium* Wilt Resistant,

PMR=Powdery Mildew Resistant,

PMT=Powdery Mildew Tolerant,

PR=*Phytophthora* Resistant,

ZYMVR=Zucchini Yellow Mosaic Virus Resistant.

Pumpkins			
Pumpkins, Less than 1 pound	WeeeeeOne* (PMR)	Pumpkins 10 to 20 pounds	Carbonado Gold* (PMT)
	Jill Be Little* (PMR)		Hermes* (PMT)
	Wee-B-Little*		Orange Sunrise* (PMT)
	Casperita		Secretariat* (PMR)
			HSC151 (edible seeds)
Pumpkins 1 to 3 pounds	Jack Sprat* (PMT)	Pumpkins More than 20 pounds	Cronos* (PMT)
	Baby Bear*		Kratos* (PMT)
	Little Giant* (PMT)		Gladiator* (PMT)
	Touch of Autumn* (PMT)		Aladdin* (PMT)
Pumpkins 2 to 6 pounds	Prankster* (PMT)		Gold Medal*
	Cannonball* (hard shell)		Rhea* (PMT)
	Iron Man* (FR, PR, PMT) (hard shell)		Solid Gold*
	Field Trip*(PMT)		Captain Jack*
	Orange Smoothie* (hard shell)	Pumpkins More than 50 pounds	Atlantic Giant
	Hybrid Pam*		Prizewinner
	Fall Splendor Plus*(PMT)	Pumpkins, Ornamental	Knucklehead*
	Mystic Plus* (PMT) (5-6 lb, plant at closer spacing to reduce size)		Goosebumps II*
	Small Sugar (BRT)	Pumpkins, Processing	Neck Pumpkin Types
	Naked Bear (ornamental, edible seeds)		Autumn Buckskin*
		Dickenson Field Types	

Winter Squash			
Winter Squash Acorn Type	Table Ace*	Winter Squash Hubbard Type	Green Hubbard
	Taybelle* (semi bush, PMT)		Golden Hubbard
	Table Gold		New England Blue Hubbard
	Table Star* (PMT)		Blue Ballet
	Autumn Delight* (PMT)		Other Hubbard Types
	Celebration* (PMT, specialty)		Boston Marrow Types
Winter Squash Butternut Type	Early Butternut*	Spaghetti Squash	Pinnacle
	Prism* (restricted vine)		Primavera*
	Metro* (restricted vine, PMR)		Vegetable Spaghetti
	Quantum*	Processing Squash	Atlas*
Waltham Butternut	Genesis*		
Winter Squash Buttercup Type	Sunshine*(orange)		Other Butternut Types
	Buttercup		
	Sweet Mama*		
	Bonbon*		

F. Pumpkins and Winter Squash

Recommended Nutrients Based on Soil Tests

In addition to using the table below, check the suggestions on rate, timing, and placement of nutrients in your soil test report and chapter B Soil and Nutrient Management. Your state's soil test report recommendations and/or your farm's nutrient management plan supersede recommendations found below.

Pumpkins and Winter Squash ¹	N (lb/A)	Soil Phosphorus Level				Soil Potassium Level				Nutrient Timing and Method
		Low	Med	High (Opt)	Very High	Low	Med	High (Opt)	Very High	
		P ₂ O ₅ (lb/A)				K ₂ O (lb/A)				
	50-100	150	100	50	0 ²	200	150	100	0 ²	Total nutrient recommended
	25-50	150	100	50	0 ²	200	150	100	0 ²	Broadcast and disk-in
	25-50	0	0	0	0	0	0	0	0	Sidedress when vines start to run

For crops grown on plastic mulch, fertilization rates are based on a standard row spacing of 6 ft.

¹Apply 20-30 lb/A of sulfur (S) for most soils.

²In VA, crop replacement values of 25 lb/A of P₂O₅ and 50 lb/A of K₂O are recommended on soils testing Very High.

Seed Treatment

Check if seed has been treated with an insecticide and fungicide. See Disease Control below.

Planting and Spacing

Seed or transplant in the field between June 15 and July 5 in cooler areas, and between May 15 and July 15 in warmer, southern areas. Base plant spacing on vine habit and average fruit size of the variety. **Note.** Fruit size may be decreased at closer spacings.

Small vine/bush with fruit less than 8 lb: Rows 5-6 ft apart with 2 ft between plants in the row.

Large/medium vine with fruit 8-15 lb: Rows 6-7.5 ft apart with 3-4 ft between plants in the row.

Large vine with fruit 12 to 25 lb: Rows 7.5-9 ft apart with 4 ft between plants in the row.

Large vine with fruit over 30 lb: Rows 10-12 ft apart with 5-6 ft between plants in the row.

Conservation Tillage (No-Till) Pumpkins

Seed or transplanted no-till pumpkins planted into small grain cover crop or stubble, hairy vetch, or fallow ground has produced commercially acceptable yields. A cover crop on the soil surface will reduce dirty pumpkins at harvest, provide some weed suppression, and minimize fruit rot by creating a barrier between pumpkins and the soil. Since cultivation is not an option in a no-till planting system and few postemergence herbicides are available to control escaped weeds, choose fields carefully for no-till production. The performance of residual preemergence herbicides depends on rainfall or overhead irrigation for activation. Moisture for activation is more critical in no-till fields consisting of a trash or straw layer. Postemergence control of grasses can be accomplished with Poast or Select. Sandea is labeled for postemergence control of yellow nutsedge and certain annual broadleaf weeds. Sandea can cause pumpkin stunting, see comments section below for more information. Sandea is an ALS inhibitor (Group 2) and is at high risk for weed resistance development. **Not recommended in NJ due to the high risk of weed resistance development and the lack of postemergence control options for certain pigweed species, common lambsquarters, annual morningglory, Eastern black nightshade, or any ALS resistant weed.**

Cover Crop Establishment and Weed Management

Small grain stubble provides an ideal crop-mulch for pumpkins.

Make sure the combine distributes straw uniformly. No other manipulation of the residue is required before planting pumpkins. An alternative crop-mulch is hairy vetch; seed in the fall 3-4 weeks before the average frost date at the rate of 20-25 lb/A with a grain drill or broadcast spreader. On sloping ground, mix a winter-killed variety of spring oats (0.5 bushel/A) with the vetch to decrease the time required for ground cover to reduce soil erosion. Adjust soil pH before vetch is seeded as tillage will not be performed before pumpkin planting. Application of P and K before seeding vetch is optional, depending on soil test results.

Soil moisture prior to planting is a critical factor for successful establishment of pumpkins.

The living, hairy vetch cover crop may remove soil moisture and prevent pumpkin germination and growth. If irrigation is not available, kill the vetch 10-14 days prior to planting in order for rainfall to provide adequate soil moisture for seeding or transplanting. If rainfall is excessive, hairy vetch may remove water to facilitate timely planting. Irrigation will eliminate the concerns about soil moisture for pumpkin seeding and germination.

Termination of the Cover Crop

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
9	Roundup PowerMax 4.5L "Generic" glyphosate 3L	16 to 32 fl oz/A 24 to 48 fl oz/A	glyphosate	0.75 to 1.13 lb acid equivalent/A	--	4
<p>-Allow at least 5-7 days between application and planting. -Some glyphosate formulations may require an adjuvant, refer to the label. -Glyphosate is not very effective for control of legumes (hairy vetch or crimson clover); glyphosate is preferred for the control for grasses. -Glyphosate-resistant horseweed is widespread in the region and will not be controlled with glyphosate. -Repeat applications are allowed, with maximum application of 5.3 qt/A per year.</p>						
22	Gramoxone SL 2.0*	2.4 to 4 pt/A	paraquat	0.6 to 1 lb/A	--	24
<p>-Apply before planting, a second application maybe required for complete control. -Always include an adjuvant (nonionic surfactant or crop oil concentrate). -Tank mix with appropriate herbicides for residual weed control; see Weed Control For Seeding Into Soil Without Plastic Mulch. -Paraquat may not control established grasses. Spray coverage is essential for optimum control. -See the label for additional information and warnings. Rainfastness is 30 min. A maximum of 3 applications per year are allowed. -Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

Pumpkin Planting

See the herbicide recommendations for pumpkins for further discussion. Use "no-till" corn planters equipped with coulters to cut through straw or cover crop stems killed by contact herbicides. Planters with finger pickup or air/vacuum units function well for seeding pumpkins. Plate planters may damage seed and should be evaluated carefully before use. Cole plate planters are satisfactory. A disk coulters on the seeding unit is essential to cut through the vetch or straw stems. Mount a 3-inch wide waffle coulters ahead of pot-transplanters to provide for effective penetration of the cover crop and plant placement.

Fertility

Hairy vetch will normally supply all the N requirements for pumpkins. However, if N deficiency symptoms appear before fruit production, topdress with 20-30 lb N/A. P and K amendments can be applied (based on soil tests) to the soil surface before planting cover crop or before planting pumpkins. When planting pumpkins into non-legume cover crops for grain stubble, apply the recommended P, K, lime, and other nutrients based on soil tests before planting. N rate recommendations may need to be increased based on fertilizer source, fertilizer application method, crop residue amount, and amount of time in a conservation tillage (no-till) production system. See section A 6. Conservation Tillage Crop Production.

Pollination (see also section A 12. Pollination).

Honey bees, squash bees, bumble bees and other wild bees are important for proper set and pollination. Populations of pollinating insects may be adversely affected by insecticides applied to flowers or weeds in bloom. Apply insecticides only in the evening hours or wait until bloom is completed before application. Check the pesticide tables below for relative toxicity to bees.

Harvest and Post-Harvest Considerations

Disease-free fruit following a regular fungicide program during crop production will minimize post-harvest fruit rots. Harvest when fruits are mature and prior to frost. Use care in handling fruit to prevent wounds. **Wounding can negate benefits from a season-long fungicide program.** Cure fruit after harvest at temperatures between 80 and 85°F (27-29°C) with a relative humidity of 75-80% for approximately 10 days. Temperatures below 50°F (10°C) cause chilling injury.

The hard-shelled squashes, such as Butternut, Delicious, Spaghetti, and the Hubbard types, can be stored at 55°F (13°C) and 50-70% relative humidity. Acorn squash will store for 5-8 weeks; pumpkins for 2-3 months and other hard-shelled squashes will store for 3 months except Hubbard types that may hold for 5-6 months. Remove squash from the field before they have chilling injury and do not allow fruits to be exposed to extended periods below 50°F (10°C). Handle fruits carefully to eliminate bruising or damage and remove stems from squash like butternuts that can damage adjacent fruit. Store winter squash in a cool, dry, well-ventilated area. The longer keeping winter squash types can be kept in saleable condition through late winter into spring (3-6 months). Research has not documented any benefit to post-harvest fruit fungicide dips.

Weed Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Herbicides

1. Identify the weeds in each field and select recommended herbicides. More information is available in the “Herbicide Effectiveness on Common Weeds in Vegetables” (Table E-3) in chapter E Pest Management.
2. Minimize herbicide resistance development. Identify the herbicide mode of action group number and follow recommended good management practices; **bolded group numbers in tables below are herbicides at higher risk for selecting resistant weed populations.** Include non-chemical weed control whenever possible.

Labeled Application Sites for Pumpkins

Herbicide (*=Restricted Use)	HRAC group number	Plastic mulch production					Bareground production		
		Soil-Applied		Postemergence			Soil-applied	POST	Post-harvest
		Under Plastic	Row Middles	Over Plastic	Row Middles	Post-Harvest			
Sandea	2		YES		YES		YES	YES	
Curbit	3		YES				YES		
Prefar	8	YES	YES				YES		
Command	13		YES				YES		
Strategy			YES				YES		
Reflex ¹	14		YES		YES		YES ²		
Dual	15		YES				YES ²		
Select	1			YES				YES	
Select Max	1			YES				YES	
Poast	1			YES				YES	
Gramoxone* ¹	22				YES	YES	YES ³		YES

¹ Special Local Needs Label 24(c), be sure it is registered for the specific state and for the intended use.

² Dual and Reflex are labeled for bareground only if the spray is directed to the row middles.

³ Apply pre-plant or after seeding but before crop emergence.

1. Soil Applied

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p>-Plasticulture row middles application only: apply before or after weed emergence; apply as a shielded application to avoid contact with the crop. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v or include a non-selective herbicide.</p> <p>-Bareground: apply broadcast after seeding but before crop emergence or no sooner than 7 days before transplanting.</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf weeds. Sandea provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field. -Do not apply Sandea to crops treated with a soil applied organophosphate insecticide or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Maximum number of Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season.</p>						
3	Curbit 3EC	1 to 3 pt/A	ethalfluralin	0.38 to 1.13 lb/A	--	24
<p>-Plasticulture: row middles only: apply as a banded spray after crop emergence or transplanting. Do not soil incorporate.</p> <p>-Bareground: apply broadcast after direct-seeding but prior to crop emergence; do not use on transplanted pumpkins.</p> <p>-Controls annual grasses and certain annual broadleaf weeds, including carpetweed and pigweed sp.</p> <p>-Use lower rate for coarse-textured soils or soils with low organic matter.</p> <p>-Where overhead irrigation is available, activate Curbit with 0.5 inch of irrigation within 2 days after application; if no irrigation or rainfall occurs within 5 days of application, activity of Curbit can be reduced.</p> <p>-Available as a pre-mix herbicide Strategy. Strategy at 3 pt/A= Curbit at 26 fl oz (0.6 lb ai) and Command at 8 fl oz (0.188 lb ai)</p> <p>-Maximum applications per season: not specified</p>						
3 + 13	Strategy 2.1SC	1.5 to 6 pt/A	ethalfluralin plus clomazone	0.39 to 1.58 lb/A	45	24
<p>-Plasticulture: row middles application only.</p> <p>-Bareground: apply broadcast just before planting or after planting but before crop emergence.</p> <p>-Strategy is a prepackage mixture of Curbit 3EC and Command 3ME. Refer to individual products for comments.</p> <p>-Clomazone spray or vapor drift may injure susceptible crops and other vegetation, refer to Command 3ME for comments.</p>						

1. Soil Applied - Strategy continued next page

1. Soil Applied - Strategy continued

<p>-Do not apply prior to planting crop. Do not soil incorporate. -Certain crop varieties may have the potential for injury or loss with this product. Consult qualified crop advisors for information pertaining to varieties in your area. -Maximum applications per season: not specified.</p>						
8	Prefar 4E	5 to 6 qt/A	bensulide	5 to 6 lb/A	--	12
<p>-Plasticulture: under plastic: apply in a band under the plastic, immediately before laying the mulch. Allow 7 day before making transplant holes to allow condensation to incorporate the herbicide. Plasticulture: row middles application is labeled. -Bareground: apply preemergence or pre-plant incorporated. -Preemergence applications should be followed by irrigation within 36 h (apply enough water to wet the soil at least 2 to 4 inches deep). Pre-plant incorporated applications should be incorporated 1 to 2 inches deep (deeper than 2 inches will result in reduced weed control). -Provides control/suppression of some annual grass weeds and some broadleaves including pigweeds, purslane, and lambsquarters. -Do not apply within 45 days of harvesting squash. -Do not apply more than 6 lb ai/A per season.</p>						
13	Command 3ME	0.67 to 2 pt/A	clomazone	0.25 to 0.75 lb/A	45	12
<p>-Labeled for winter squash and processing pumpkins; not labeled for jack-o-lantern pumpkins. -Plasticulture: row middles application only. -Bareground: apply broadcast just before planting but before crop emergence, or just before transplanting. Use the lower rate when used on coarse-textured soils low in organic matter, when weed pressure is light, or to minimize herbicide carryover that could affect subsequent crops. -Controls annual grasses and many broadleaf weeds including common lambsquarters, velvetleaf, spurred anoda, and jimsonweed. Carpetweed, morningglory sp., pigweed sp., and yellow nutsedge will not be controlled. Higher rates will improve control (or expand number of species controlled) such as common cocklebur, common ragweed, or jimsonweed (refer to label for specific weeds and rates). -WARNINGS: Command spray or vapor drift may injure sensitive crops and other vegetation up to several hundred yards from the point of application. Do not apply adjacent to sensitive crops (see label) or vegetation, or under unfavorable wind or weather conditions. Command may limit subsequent cropping options, see the label. -Available as a pre-mix herbicide Strategy: Strategy at 3 pt/A= Command at 8 fl oz (0.188 lb ai) and Curbit at 26 fl oz (0.6 lb ai) -Maximum number of Command applications per year: 1.</p>						
14	Reflex 2SL	Rates vary, refer to the specific label	fomesafen	0.13 to 0.38 lb/A	32	24
<p>-For pumpkins ONLY. A Special Local Needs Label 24(c) has been approved for the use of Reflex 2SL to control weeds in pumpkins in DE, NJ, PA (expires 12/31/2025 in DE, 12/31/2022 in NJ, and 12/31/2023 in PA). The use of this product is legal ONLY if a waiver of liability has been completed (see https://www.syngenta-us.com/labels/indemnified-label-login). -Rates differ by states, soil types, and planting method. Rates as low as 10 fl oz/A can cause injury on coarse-textured soils. -Plasticulture: row middles application only. -Bareground: apply broadcast within 24 h after direct-seeding and follow with 0.2 to 0.5 inches of overhead irrigation at least 36 h before pumpkin begin to crack through the soil. For transplants, apply Reflex and then irrigate with 0.2 to 0.5 inches of water and then transplant. Do not prepare transplant holes until after Reflex application and irrigation. -Foliar application of Reflex will severely damage or kill pumpkin. The potential of crop injury is greater on lighter textured soils combined with intensive irrigation programs or high amounts of rainfall, therefore, adjust rates accordingly. -Reflex provides both residual and postemergence control of susceptible weed species. Effective postemergence control requires an adjuvant. Pumpkin varieties may vary in their response to Reflex. Treat small acreages first to determine tolerance, especially when applying to a new variety. -Reflex rates lower than 16 fl oz/A may not provide full-season control and should be used with other herbicides and/or other methods of weed control. -Consider rotational crops when applying fomesafen. If crop is replanted, do not re-apply Reflex. Refer to 24(c) label for specifics on rotational restrictions. -Maximum for Reflex application in DE, MD, NJ, PA, and VA: 24 fl oz/A IN ALTERNATE YEARS.</p>						
15	Dual Magnum 7.62E	1 to 1.33 pt/A	s-metolachlor	0.95 to 1.27 lb/A	30	24
<p>-For pumpkins ONLY. Plasticulture: row middles application only. Bareground: apply as an inter-row or inter-hill spray, leaving 1 ft of untreated area over the row. Do not use as an over the top application. Do not soil incorporate. -Suppresses or controls annual grasses, yellow nutsedge, and certain annual broadleaf weeds including nightshade species. -Dual Magnum will not control emerged weeds. Cultivate and/or hoe or tank mix with Gramoxone to control emerged weeds before treatment. -Use the lower rate on fields with coarse-textured soils low in organic matter. Use the higher rates on fields with fine-textured soil and those with high organic matter. Maximum applications per season: not specified.</p>						

2. Postemergence

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
1	Select 2EC	6 to 8 fl oz/A	clethodim	0.07 to 0.125 lb/A	14	24
	Select Max 0.97EC	9 to 16 fl oz/A				
	Poast 1.5EC	1 to 1.5 pt/A	sethoxydim	0.19 to 0.28 lb/A	14	12
<p>-Select 2EC: use crop oil concentrate (COC) at 1% v/v (1 gal/100 gal of spray solution). Select Max: use nonionic surfactant (NIS) at 0.25% v/v (1 qt/100 gal of spray solution). Poast: use COC at 1.0% v/v.</p>						

2. Postemergence Select, Select Max, Poast - continued next page

F. Pumpkins and Winter Squash

2. Postemergence Select, Select Max, Poast - continued

<p>-The use of COC may increase the risk of crop injury when hot or humid conditions prevail. To reduce the risk of crop injury, omit additives or switch to NIS when grasses are small and soil moisture is adequate.</p> <p>-Use lower labeled rates for annual grass control and higher labeled rates for perennial grass control.</p> <p>-Yellow nutsedge, wild onion, wild garlic, and broadleaf weeds will not be controlled.</p> <p>-Controls many annual and certain perennial grasses, including annual bluegrass, but Poast is preferred for goosegrass control. For best results, treat annual grasses when they are actively growing and before tillers are present. Control may be reduced if grasses are large or under hot or dry weather conditions.</p> <p>-Repeated applications may be necessary to control certain perennial grasses. If repeat applications are necessary, allow 14 days between applications.</p> <p>-Rainfastness is 1 h.</p> <p>-Do not tank mix with or apply within 2 to 3 days of any other pesticide, unless labeled, as this may increase the risk of crop injury or reduce the control of grasses. Do not apply more than 8 fl oz of Select 2EC in a single application and do not exceed 32 fl oz/A for the season; do not apply more than 16 fl oz of Select Max in a single application and do not exceed 64 fl oz/A for the season.</p> <p>-Do not apply more than 1.5 pt/A Poast in a single application and do not exceed 3 pt/A for the season.</p>						
2	Sandea 75DF	0.5 to 1 oz/A	halosulfuron	0.023 to 0.047 lb/A	30	12
<p>-Plasticulture: row middles application only.</p> <p>-Bareground: broadcast for bareground. Apply Sandea after the crop has at least 3 to 5 true leaves but before first female flowers appear and no sooner than 14 days after transplanting. If weeds have emerged, use a non-ionic surfactant at 0.25% v/v (1 qt/100 gal).</p> <p>-Suppresses or controls yellow nutsedge and certain broadleaf; control of weeds taller than 3 inches may not be adequate. Sandea will not control common lambsquarters or eastern black nightshade if applied postemergence; for row middle application, tank mix with a non-selective herbicide to increase spectrum of control.</p> <p>-Sandea provides both residual and postemergence control of susceptible weed species.</p> <p>-Sandea is an ALS inhibiting herbicide and resistant weed populations are common in the region. Do not use Group 2 herbicides repeatedly in the same field.</p> <p>-Do not apply Sandea to crops treated with a soil applied organophosphate insecticide, or use a foliar applied organophosphate insecticide within 21 days before or 7 days after a Sandea application.</p> <p>-Rainfastness is 4 h. Maximum number of Sandea applications per year is 2 and do not exceed 2 oz/A during the crop season</p>						
22	Gramoxone SL 2.0* Gramoxone SL 3.0*	1.95 pt/A 1.3 pt/A	paraquat	0.49 lb/A	14	24
<p>-A Supplemental Label has been approved for the use of Gramoxone 2SL or 3SL for postemergence weed control in DE, MD, NJ, PA, and VA. Row middles as a shielded application. Apply as a directed spray in a minimum of 20 gal spray mix/A to control emerged weeds between the rows after crop establishment. Include a nonionic surfactant at 0.25% v/v.</p> <p>-Use shields or hoods to prevent spray contact with the crop and low spray pressure (maximum of 30 psi) to reduce small droplets that are prone to drift. See the label for additional information and warnings.</p> <p>-Rainfastness is 30 min. A maximum of 3 applications per year are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

3. Postharvest

Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient	Active Ingredient Rate	PHI (d)	REI (h)
22	Gramoxone SL 2.0* Gramoxone SL 3.0*	2.25 to 3 pt/A 1.5 to 2 pt/A	paraquat	0.56 to 0.75 lb/A	--	24
<p>-A Special Local Needs Label 24(c) has been approved for Gramoxone SL 2.0 in VA (expires 12/31/2022) and a Supplemental Label in DE for the use of both Gramoxone formulations for postharvest application to desiccate the crop. Apply after the last harvest for bareground or plasticulture.</p> <p>-Always include an adjuvant. Spray coverage is essential for optimum effectiveness.</p> <p>-See the label for additional information and warnings.</p> <p>-Rainfastness 30 min. A maximum of 2 applications for crop desiccation are allowed.</p> <p>-Restricted-use pesticide. Only certified applicators, who successfully complete the paraquat-specific training, can mix, load or apply paraquat. Application of paraquat "under the direct supervision" of a certified applicator is no longer allowed. Required training link (http://usparaquattraining.com); certified applicators must repeat training every three years.</p>						

4. Other Labeled Herbicides

These products are labeled but limited local data are available; and/or are labeled but not recommended in our region due to potential crop injury concerns.

Group	Product Name (*=Restricted Use)	Active Ingredient
14	Aim	carfentrazone
14	Valkos 51 WDG	flumioxazin
14	Vida	pyraflufen

Insect Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F. Recommended Insecticides

Seed and At-Plant Treatments for Seedcorn Maggot

Farmore DI-400 as a commercially applied seed treatment which contains thiamethoxam (Group 4A).

Athena* (bifenthrin + avermectin B1, Group 3A + 6) at planting at 8.5 to 17 fl oz/A.

Verimark (cyantranilprole, Group 28) applied no earlier than 72 hours prior to planting, at 10-13.5 oz/A using in-furrow spray, transplant tray drench, transplant water treatment, hill drench, or surface band.

Note:

The use of neonicotinoid insecticides (Group 4A) at planting may help reduce seedcorn maggot populations. See also Maggots in section E 3.1. Soil Pests - Detection and Control.

Aphids

Note: Aphids transmit Mosaic Virus.

Apply one of the following formulations: Note: Thorough spray coverage beneath leaves is important. Treat seedlings every 5-7 days, or as needed.						
Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	3.0 fl oz/A	afidopyropen	0	12	L
21A	Torac	17.0 to 21.0 fl oz/A	tolfenpyrad	1	12	H
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	Soil, at planting: 10 to 13.5 fl oz/A Drip chemigation: 10 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro*	10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H
29	Beleaf 50SG	Foliar: 2.0 to 2.8 oz/A Drip: 2.8 to 4.28 oz/A	flicicamid	0	12	L

Armyworms and Cabbage Loopers

Apply one of the following formulations:						
Group	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Proclaim 5SG*	3.0 to 4.8 oz/A	emamectin benzoate	7	12	H
11A	Dipel DF, others (OMRI)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis kurstaki</i>	0	4	N
11A	XenTari (OMRI) (armyworms)	0.5 to 2.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
11A	XenTari (OMRI) (cabbage loopers)	0.5 to 1.0 lb/A	<i>Bacillus thuringiensis aizawai</i>	0	4	N
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	3.5 to 7.5 fl oz/A	chlordantraniliprole - soil and foliar	1	4	L
28	Exirel (armyworms)	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Exirel (cabbage loopers)	10.0 to 17.0 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Voliam Flexi (cabbage looper only)	4.0 to 7.0 oz/A	thiamethoxam + chlordantraniliprole	1	12	H
28 + 6	Minecto Pro*	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H

F. Pumpkins and Winter Squash

Cucumber Beetles

Young plants need to be protected from cucumber beetle feeding as the beetles can transmit the causal agent of bacterial wilt. Cucumber beetles also cause direct damage to pumpkin and winter squash rinds. Management of adult cucumber beetles early in the season may help reduce damage to rinds later in the season. Seeds pretreated with a neonicotinoid seed treatment such as Farmore DI400 should provide up to 14 days of control of cucumber beetle. Otherwise, apply one of the following formulations:

Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
28	Exirel	20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	Soil, at planting: 13.5 fl oz/A Drip chemigation: 10 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H

Cutworms See also section E 3.1. Soil Pests - Detection and Control.

Apply one of the following formulations:						
Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					

Leafminers

Apply one of the following formulations:						
Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Agri-Mek SC*	1.75 to 3.5 fl oz/A	abamectin	7	12	H
17	Trigard 75WSP	2.66 oz/A	cyromazine	0	12	H
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - soil and foliar	1	4	L
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 6	Minecto Pro*	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H

Melonworms and Pickleworms

Apply one of the following formulations. When using foliar materials make one treatment prior to fruit set, and then treat weekly. For soil or drip applications check the label for instructions on treatment frequency.						
Group	Product Name (*= Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	0.5 to 1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Proclaim 5SG*	3.0 to 4.8 oz/A	emamectin benzoate	7	12	H
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L
22	Avaunt 30WDG, Avaunt eVo	2.5 to 6.0 oz/A	indoxacarb	3	12	H
28	Coragen 1.67SC	5.0 to 7.5 fl oz/A	chlorantraniliprole - soil	1	4	L
28	Coragen 1.67SC	2.0 to 3.5 fl oz/A	chlorantraniliprole - foliar	1	4	L
28	Exirel	7.0 to 13.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	5.0 to 10.0 fl oz/A	cyantraniliprole	1	4	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclaniliprole	1	4	H
28 + 4A	Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole	30	12	H
28 + 4A	Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole	1	12	H
28 + 6	Minecto Pro*	5.5 to 10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H

Mites

Mite infestations generally begin around field margins and grassy areas. **DO NOT** mow or maintain these areas after midsummer to prevent mites from moving into the crop. Localized infestations can be spot-treated. Begin treatment when 10-15% of the crown leaves are infested early in the season.

Apply one of the following formulations. Note: Continuous use of carbaryl or pyrethroids may result in mite outbreaks. Addition of crop oils or organosilicon spray additives will increase miticide effectiveness.						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
6	Agri-Mek SC*	1.75 to 3.5 fl oz/A	abamectin	7	12	H
6 + 3A	Athena*	13.5 to 17 fl oz/A	avermectin B1 + bifenthrin	7	12	H
10B	Zeal Miticide	2.0 to 3.0 oz/A	etoxazole	7	12	L
20B	Kanemite 15SC	31.0 fl oz/A	acequinocyl	1	12	L
21A	Magister SC	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28 + 6	Minecto Pro*	5.5 to 10.0 fl oz/A	cyantranilprole + abamectin	7	12	H
20D	Acramite 50WS	0.75 to 1.0 lb/A	bifenazate	3	12	M

Rindworms

For Lepidopteran Rindworms, apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	4.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	5.0 to 10.0 fl oz/A	spinetoram	3	4	M
6	Proclaim 5SG*	3.0 to 4.8 oz/A	emamectin benzoate	7	12	H
18	Intrepid 2F	4.0 to 10.0 fl oz/A	methoxyfenozide	3	4	L

Squash Bugs

Begin treatments if more than one egg mass per plant is present. Sprays should target nymphal stages.

Apply one of the following formulations: Note: Under-leaf spray coverage is essential.						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Sevin XLR Plus	1.0 qt/A	carbaryl	3	12	H
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	10.5 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M

Squash Vine Borers

When vines begin to run, apply to bases of plants 4 times at 7-day intervals. Pheromone traps for squash vine borer are commercially available. These traps can be used to indicate when moth activity begins. Note: Use of spinosad or spinetoram for Cabbage Looper control will reduce squash vine borer populations.

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					

Thrips

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
3A ¹	Pyrethroid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4A ²	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
5	Entrust SC (OMRI)	6.0 to 8.0 fl oz/A	spinosad	3	4	M
5	Radiant SC	6.0 to 10.0 fl oz/A	spinetoram	3	4	M
21A	Torac	21.0 fl oz/A	tolfenpyrad	1	12	H
28	Harvanta 50SL	10.9 to 16.4 fl oz/A	cyclanilprole	1	4	H
29	Beleaf 50SG	Foliar: 2.0 to 2.8 oz/A Drip: 2.8 to 4.28 oz/A	flonicamid	0	12	L

¹Resistance concerns with western flower thrips ²Resistance concerns with tobacco thrips

F. Pumpkins and Winter Squash

Whiteflies

Apply one of the following formulations:						
Group	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
4A	Neonicotinoid insecticides registered for use on Pumpkins and Winter Squash: see table at the end of Insect Control.					
4D	Sivanto Prime or 200SL	7.0 to 14.0 fl oz/A	flupyradifurone - foliar	1	4	M
4D	Sivanto Prime or 200SL	21.0 to 28.0 fl oz/A	flupyradifurone - soil	21	4	M
7C	Knack	8.0 to 10.0 fl oz/A	pyriproxyfen	7	12	L
9B	Fulfill 50WDG	2.75 oz/A	pymetrozine	0	12	L
9B	PQZ	2.4 to 3.2 fl oz/A	pyrifluquinazon	1	12	L
9D	Sefina	14.0 fl oz/A	afidopyropen	0	12	L
23	Oberon 2SC	7.0 to 8.5 fl oz/A	spiromesifen	7	12	M
28	Exirel	13.5 to 20.5 fl oz/A	cyantraniliprole	1	12	H
28	Verimark	6.75 to 13.5 fl oz/A	cyantraniliprole	1	4	H
28 + 6	Minecto Pro*	10.0 fl oz/A	cyantraniliprole + abamectin	7	12	H
29	Beleaf 50SG	Foliar: 2.0 to 2.8 oz/A Drip: 2.8 to 4.28 oz/A	flonicamid	0	12	L

Group 3A Pyrethroid Insecticides Registered for Use on Pumpkins and Winter Squash						
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):						
Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR	
Asana XL*	5.8 to 9.6 fl oz/A	esfenvalerate	3	12	H	
Baythroid XL*	0.8 to 2.8 fl oz/A	beta-cyfluthrin	0	12	H	
Brigade 2EC*, others	2.6 to 6.4 fl oz/A	bifenthrin	3	12	H	
Danitol 2.4EC*	10.67 to 16.0 fl oz/A	fenpropathrin	7	24	H	
Declare*	1.02 to 1.54 fl oz/A	gamma-cyhalothrin	1	24	H	
Hero EW*	4.0 to 10.3 fl oz/A	zeta-cypermethrin + bifenthrin	3	12	H	
Lambda-Cy 1EC*, others	2.56 to 3.84 fl oz/A	lambda-cyhalothrin	1	24	H	
Mustang Maxx*	1.28 to 4.0 fl oz/A	zeta-cypermethrin	1	12	H	
Permethrin 3.2EC*, others	4.0 to 8.0 fl oz/A	permethrin	0	12	H	
Tombstone*, others	0.8 to 2.8 fl oz/A	cyfluthrin	0	12	H	
Warrior II*	1.28 to 1.92 fl oz/A	lambda-cyhalothrin	1	24	H	
Combo products containing a pyrethroid						
Athena*	7.0 to 17.0 fl oz/A	bifenthrin + avermectin B1 (Group 6)	7	12	H	
Besiege*	6.0 to 9.0 fl oz/A	lambda-cyhalothrin + chlorantraniliprole (Group 28)	1	24	H	
Endigo ZC*	4.0 to 4.5 fl oz/A	lambda-cyhalothrin + thiamethoxam (Group 4A)	1	24	H	
Gladiator*	19.0 fl oz/A	zeta-cypermethrin + abamectin (Group 6)	7	12	H	
Savoy EC*	6.0 to 12.9 fl oz/A	bifenthrin + acetamiprid (Group 4A)	7	12	H	

Group 4A Neonicotinoid Insecticides Registered for Use on Pumpkins and Winter Squash						
Apply one of the following formulations (check if the product label lists the insect you intend to spray; the label is the law):						
Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR	
Admire Pro	7.0 to 10.5 fl oz/A	imidacloprid - soil	21	12	H	
Assail 30SG	2.5 to 5.3 oz/A	acetamiprid	0	12	M	
Belay 2.13SC	9.0 to 12.0 fl oz/A	clothianidin - soil/drip	21	12	H	
Belay 2.13SC	3.0 to 4.0 fl oz/A	clothianidin - foliar (note: phi: do not make application after 4 th true leaf has unfolded)	see note	12	H	
Actara 25WDG	1.5 to 5.5 oz/A	thiamethoxam	0	12	H	
Platinum 75SG	1.66 to 3.67 oz/A	thiamethoxam	30	12	H	
Scorpion 35SL	9.0 to 10.5 fl oz/A	dinotefuran - soil/drip	21	12	H	
Scorpion 35SL	2.0 to 7.0 fl oz/A	dinotefuran - foliar	1	12	H	
Venom 70SG	5.0 to 7.5 oz/A	dinotefuran - soil/drip	21	12	H	
Venom 70SG	1.0 to 4.0 oz/A	dinotefuran - foliar	1	12	H	
Combo products containing a neonicotinoid						
Durivo	10.0 to 13.0 fl oz/A	thiamethoxam + chlorantraniliprole (Group 28)	30	12	H	
Endigo ZC*	4.0 to 4.5 fl oz/A	thiamethoxam + lambda-cyhalothrin (Group 3A)	1	24	H	
Savoy EC*	6.0 to 12.9 fl oz/A	acetamiprid + bifenthrin (Group 3A)	7	12	H	
Voliam Flexi	4.0 to 7.0 oz/A	thiamethoxam + chlorantraniliprole (Group 28)	1	12	H	

Disease Control

THE LABEL IS THE LAW-see the Pesticide Use Disclaimer on the first page of chapter F.
Recommended Fungicides

Nematodes See also sections E 1.5. Soil Fumigation and E 1.6. Nematode Control. Use fumigants listed in section E 1.5., or nematicides listed below. Consult the label.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
1A	Vydate L*	1.0 to 2.0 gal/A incorporate into top 2-4 inches of soil, <i>OR</i> 2.0 to 4.0 pt/A apply 2 w after planting and repeat 2-3 w later.	oxamyl	1	48	H
7	Velum Prime 4.16SC	6.5 to 6.84 fl oz/A	fluopyram	0	12	--
--	Nimitz 4EC	3.5 to 5.0 pt/A incorporate or drip-apply 7 d before planting	fluensulfone	n/a	12	N

Seed Treatment

Check with your seed company if seed has been treated with an insecticide and fungicide. If it has not been treated, use a mixture of Thiram 480DP (4.5 fl oz/100 lb seed) and an approved commercially available insecticide.

Damping-off caused by *Phytophthora*, *Pythium*, and *Rhizoctonia*

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Apply one of the following at-planting (see label for application timing, methods, and restrictions):						
Phytophthora and Pythium Root Rot						
4	Ridomil Gold 4SL ¹	1.0 to 2.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E ¹	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4	MetaStar 2E AG ¹	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
Phytophthora, Pythium, and Rhizoctonia Root Rot						
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row. Avoid direct seed contact, which may cause delayed emergence.	mefenoxam + azoxystrobin	AP	0	N
Rhizoctonia root rot						
11	azoxystrobin 2.08F	0.40 to 0.80 fl oz/1000 ft row	azoxystrobin	1	4	N
Pythium root rot only						
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or direct spray at base of plant and soil	propamocarb hydrochloride	2	12	N

¹To determine the amount of Ridomil Gold, Ultra Flourish, or MetaStar needed per acre, use the following calibration formula for changing from broadcast to band application: [Band width (ft) / row spacing (ft)] x broadcast rate (lb/A) = Amount needed lb/A.

Bacterial and Fungal Diseases

Angular Leaf Spot/Bacterial Leaf Spot

Both diseases can produce foliar symptoms that are often overlooked. Early detection is important, since control of the foliar phase can reduce infections in developing fruit. Infected fruit will become unmarketable. Both diseases are seedborne and can survive on infested debris for at least one year or until the debris decomposes. Rotate away from fields with a history of bacterial problems. Incorporate the following into a standard disease management program when leaf spot is first detected and repeat every 7 to 10 days: fixed copper at labeled rates plus mancozeb.

Anthracnose - see Gummy Stem Blight (Black Rot) and Anthracnose below.

Bacterial Wilt

Controlling striped and spotted cucumber beetles is essential for preventing bacterial wilt. See "Cucumber Beetles" in the Cucumber Insect Control section for specific recommendations. Insecticide applications made at planting may not prevent beetle damage season-long; additional foliar insecticide applications may be necessary.

Choanophora Fruit Rot

This disease occurs during warm wet weather and develops predominantly on flowers or fruit near the ground. Management is difficult because disease development is rapid and weather dependent. Fungicide sprays are not effective because flowers, which open daily, must be protected immediately. Practices that reduce soil moisture or reduce flower-soil contact, such as raised beds and plastic mulch, may be beneficial.

F. Pumpkins and Winter Squash

Downy Mildew

Scout fields for disease incidence on a regular basis. Begin targeted sprays when Downy Mildew is predicted for the region. For current status of the disease, check the Cucurbit Downy Mildew Forecasting website at <https://cdm.ipmpipe.org>. Strains of Downy Mildew that infect one cucurbit crop may not affect pumpkin or winter squash. Unnecessary fungicide application can be avoided by not spraying until disease is predicted in the region on watermelon. Preventative applications are much more effective than applications made after disease is detected. Materials with different modes of action (FRAC codes) should always be alternated to reduce the chances for fungicide resistance development.

Code	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Sprays should be applied on a 7-d schedule when disease is forecast or present in the region. Under severe disease conditions and conducive weather, spray interval may be reduced IF the label allows. TANK-MIX one of these products WITH a protectant fungicide such as chlorothalonil 6F or Gavel 75DF:						
49 + 40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
21	Ranman 400SC	2.10 to 2.75 fl oz/A (do not apply with copper; see label for details) ¹	cyazofamid	0	12	L
Other materials for use in rotations as tank mix partners with a protectant:						
43	Presidio 4SC	3.0 to 4.0 fl oz/A	fluopicolide	2	12	L
28	Previcur Flex 6F	1.2 pt/A	propamocarb hydrochloride	2	12	N
40 + 45	Zampro 525SC	14.0 fl oz/A	dimethomorph + ametoctradin	0	12	--
22	Elumin 4SC	8.0 fl oz/A	ethaboxam	2	12	--
M03+22	Gavel 75DF	1.5 to 2.0 lb/A contains protectant	mancozeb + zoxamide	5	48	--
M05+22	Zing! 4.9SC	36 fl oz/A contains protectant	chlorothalonil + zoxamide	0	12	N
M05+27	Ariston 42SC	1.9 to 3.0 pt/A contains protectant	chlorothalonil + cymoxanil	3	12	--
11 + 27	Tanos 50DF	8.0 oz/A	famoxadone + cymoxanil	3	12	--
27	Curzate 60DF	3.2 to 5.0 oz/A	cymoxanil	3	12	N
29	Omega 500F	12.0 to 24.0 fl oz/A	fluazinam	7	12	N
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N

¹Ranman should be tank mixed with an organosilicone surfactant when disease is severe, or a non-ionic surfactant or blend of organosilicone and non-ionic surfactant disease is moderate or light.

Fusarium Fruit Rot

This disease is especially destructive in fields where pumpkins are grown every year. Once the pathogen is established in a field, loss can be significant. Fruit Rot is caused by several *Fusarium* spp., and fungicide applications are not effective. Hard rind cultivars are less susceptible to *Fusarium* Fruit Rot than other cultivars. Production of pumpkin on a no-till cover crop mulch layer such as winter rye plus hairy vetch has been shown to help reduce disease incidence. Greater disease reductions will occur when the mulch layer is thicker.

Gummy Stem Blight (Black Rot) and Anthracnose

Rotate crops to allow at least 2 years between cucurbit plantings. Pumpkin cv. 'Small Sugar' appears to be the least affected by Black Rot.

Code	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Fungicides with a high-risk for resistance development, such as FRAC code 11 fungicides (Cabrio, Pristine and Quadris), should be tank-mixed with a protectant fungicide. Use at least the minimum labeled rate of each fungicide in the tank-mix. Do not apply FRAC code 11 fungicides more than 4 times total per season. If resistance to FRAC code 11 fungicides exists in the area, use fungicides from a different FRAC code. Begin the following fungicide program when fruit start to form: ALTERNATE:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A (use low rate early in season)	chlorothalonil	0	12	N
WITH one of the following:						
3	tebuconazole 3.6F	8.0 fl oz/A	tebuconazole	7	12	N
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	--
3	Rhyme 2.08SC	5.0 to 7.0 fl oz/A	flutriafol	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	7	12	--
3 + 7	Luna Experience 3.34SC ¹	10.0 to 17.0 fl oz/A	tebuconazole + fluopyram	7	12	--
7 + 12	Miravis Prime	9.2 to 11.4 fl oz/A	pydiflumetofen + fludioxonil	1	12	--

Gummy Stem Blight (Black Rot) and Anthracnose - continued next page

Gummy Stem Blight (Black Rot) and Anthracnose - continued

9 + 12	Switch 62.5WG	11.0 to 14.0 oz/A	cyprodinil + fludioxonil	1	12	L
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
7 + 11	Merivon 2.09SC ²	5.5 fl oz/A	fluxapyroxad + pyraclostrobin	0	12	N
7 + 11	Pristine 38WG ²	12.5 to 18.5 oz/A	boscalid + pyraclostrobin	0	12	--
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N

**Maintain fungicide schedule until harvest (see “Harvest and Post-Harvest Considerations” section above).
Fungicide application for Black Rot control will help maintain “handles” on the fruit.
Harvest carefully because wounding can negate benefits from a season-long fungicide program.**

¹A mild yellowing on leaf margins is sometimes seen following application of Luna Experience in cucurbits. ²Tank mixes of additives, adjuvants, and/or other products may result in crop injury.

Phytophthora Crown and Fruit Rot

Multiple practices should be used to minimize the occurrence of this disease. Rotate away from susceptible crops (such as peppers, eggplants, tomatoes, lima and snap beans, and other cucurbits) for as long as possible. Pre-plant fumigants will also suppress disease. Fields should be adequately drained to ensure that water does not accumulate around the base of the plant. Once the canopy closes, subsoil between the rows to allow for faster drainage following rainfall. Materials with different modes of action (*i.e.*, FRAC codes) should always be alternated to reduce the chances for fungicide resistance development. Apply fungicides when conditions are favorable for disease development. Fruit are susceptible at all growth stages and must be protected season-long.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Apply one of the following formulations pre-plant for early season control:						
4	MetaStar 2E AG	4.0 to 8.0 pt/A	metalaxyl	AP	48	N
4	Ridomil Gold 4SL	1.0 to 2.0 pt/A	mefenoxam	5	48	N
4	Ultra Flourish 2E	2.0 to 4.0 pt/A	mefenoxam	5	48	N
4 + 11	Uniform 3.66SE	0.34 fl oz/1000 ft row	mefenoxam + azoxystrobin	AP	0	N
28	Previcur Flex 6F	1.2 pt/A in transplant water, drip irrigation, or spray directed to the base of the plants and soil.	propamocarb hydrochloride	2	12	N
49 + 4	Orondis Gold 1.67SC ¹	28.0 to 55.0 fl oz/A in furrow or by drip	oxathiapiprolin + mefenoxam	5	48	--
Apply one of the following fungicides and tank mix with fixed copper at labeled rates when conditions favor disease development (for suppression only):						
49 + 40	Orondis Ultra 2.33SC	5.5 to 8.0 fl oz/A	oxathiapiprolin + mandipropamid	0	4	--
49+M05	Orondis Opti	1.75 to 2.5 pt/A	oxathiapiprolin + chlorothalonil	0	12	--
21	Ranman 400SC	2.75 fl oz/A (do not apply with copper; see label for details) ²	cyazofamid	0	12	L
40 + 45	Zampro 525SC	14.0 fl oz/A	dimethomorph + ametoctradin	0	12	--
22	Elumin 4SC	8 fl oz/A	ethaboxam	2	12	--
40	Revus 2.08F	8.0 fl oz/A	mandipropamid	0	4	--
40	Forum 4.17SC	6.0 fl oz/A	dimethomorph	0	12	N
43	Presidio 4SC ³	4.0 fl oz/A	fluopicolide	2	12	L
M05+22	Zing! 4.9SC	36 fl oz/A	chlorothalonil + zoxamide	0	12	N

¹Do not follow soil applications of Orondis Gold 1.67SC with foliar applications of oxathiapiprolin-containing products. ²Ranman should be tank mixed with an organosilicone surfactant when disease is severe, or a non-ionic surfactant or blend of organosilicone and non-ionic surfactant disease is moderate or light. ³Presidio may also be applied through the drip irrigation (see supplemental label). Soil drench followed by drip application has given good results in some trials on crown rot caused by *Phytophthora capsici*.

Plectosporium Blight (Microdochium blight)

Research has shown that no-till pumpkin production may reduce disease. Rotate with crops other than cucurbits. It is important to achieve maximum foliage coverage with each fungicide application. Scout fields regularly.

Code	Product Name (* = Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Once symptoms appear on petioles or as fruit begins to form, apply one of the following and repeat every 7-10 days:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N
3 + 11	Quadris Top 1.67SC ¹	12.0 to 14.0 fl oz/A	difenoconazole + azoxystrobin	1	12	--
7 + 11	Pristine 38WG ²	18.5 oz/A	boscalid + pyraclostrobin	0	12	--

**A spray schedule that alternates Cabrio 20EG or Flint Extra 500SC with chlorothalonil will also provide control.
Note: do not apply Flint Extra 500SC near Concord grapes, see label.**

¹Do not apply near apples, see label. ²Tank mixes of additives, adjuvants, and/or other products may result in crop injury.

F. Pumpkins and Winter Squash

Powdery Mildew

Some varieties have resistance or tolerance to Powdery Mildew and should be used if possible (see table Recommended Varieties above). The fungus that causes cucurbit Powdery Mildew has developed resistance to high-risk fungicides. In the Eastern US, resistance to strobilurin (FRAC code 11) and DMI (FRAC code 3) fungicides has been reported. Proper fungicide resistance management should be followed to help delay the development of resistance and minimize control failures.

Powdery Mildew generally occurs from mid-July until the end of the season. Development on tolerant varieties will vary from year to year. Planting tolerant varieties will help delay the development of Powdery Mildew and improve the performance of fungicides. If Powdery Mildew has become well established in the mid- to late part of the season, only apply protectant fungicides such as chlorothalonil or sulfur. Make first application when Powdery Mildew is observed in the area or is detected by scouting (one lesion on the underside of 45 old leaves per acre).

Code	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
TANK MIX one of these products with a protectant such as chlorothalonil 6F 2.0 to 3.0 pt/A:						
50	Vivando 2.5SC ¹	15.4 fl oz/A	metrafenone	0	12	--
3 + 7	Luna Experience 3.34SC ²	6.0 to 17.0 fl oz/A	tebuconazole + fluopyram	7	12	--
13	Quintec 2.08SC	4.0 to 6.0 fl oz/A	quinoxifen	3	12	--
AND ALTERNATE with fungicides from different FRAC codes with a protectant such as chlorothalonil 6F 2.0 to 3.0 pt/A:						
3	tebuconazole 3.6F	4.0 to 6.0 fl oz/A	tebuconazole	7	12	N
3	Procur 480SC	4.0 to 8.0 fl oz/A	triflumizole	0	12	N
3	Proline 480SC	5.7 fl oz/A	prothioconazole	7	12	--
3	Rally 40WSP	2.5 to 5.0 oz/A	myclobutanil	0	24	N
3	Rhyme 2.08SC	5.0 to 7.0 fl oz/A	flutriafol	0	12	--
3 + 9	Inspire Super 2.82EW	16.0 to 20.0 fl oz/A	difenoconazole + cyprodinil	7	12	--
3 + 7	Aprovia Top 1.62EC	10.5 to 13.5 fl oz/A	difenoconazole + benzovindiflupyr	0	12	--
7 + 11	Pristine 38WG ³	12.5 to 18.5 oz/A	boscalid + pyraclostrobin	0	12	--
39	Magister 1.6SC ⁴	24.0 to 36.0 fl oz/A	fenazaquin	3	12	H
7 + 12	Miravis Prime	9.2 to 11/4 fl oz/A	pydiflumetofen + fludioxonil	1	12	--
P05	Regalia (OMRI)	4.0 qt/A	Extract of <i>Reynoutria sachalinensis</i>	0	4	--
OR WITH (Note: Sulfur may injure plants, especially at high temperatures. Certain varieties can be more sensitive. Consult the label for precautions).						
M02	Micronized Wettable Sulfur 80W ⁵	4.0 lb/A	sulfur	--	24	N
U06	Torino 0.85SC	3.4 fl oz/A	cyflufenamid	0	4	--

¹Do not mix Vivando with horticultural oils.

²A mild yellowing on leaf margins is sometimes seen following application of Luna Experience in cucurbits.

³Tank mixes of additives, adjuvants, and/or other products may result in crop injury.

⁴Do not make more than one application per year of Magister.

⁵Do not apply when temperature exceeds 90°F or to varieties susceptible to sulfur injury.

Scab

Select scab-resistant varieties. The fungus that causes scab typically occurs during periods of cool, wet weather when temperatures are below normal. Rotate away from fields with a history of scab for at least 2 years.

Code	Product Name (*=Restricted Use)	Product Rate	Active Ingredient(s)	PHI (d)	REI (h)	Bee TR
Begin sprays as true leaves form and repeat every 5 to 7 days:						
M05	chlorothalonil 6F	2.0 to 3.0 pt/A	chlorothalonil	0	12	N

Viruses (WMV, PRSV, ZYMV, and CMV)

The most prevalent virus in the mid-Atlantic region is WMV, followed by PRSV, ZYMV, and CMV. An easy method for mitigating potential losses are to plant varieties with resistance packages to multiple viruses whenever possible. Plant fields as far away from existing cucurbit plantings as possible to help reduce aphid transmission of viruses.

If you are having a medical emergency after using pesticides, call 911 immediately.

If you have any of the following symptoms during or shortly after using pesticides: headache, blurred vision, pinpoint pupils, weakness, nausea, cramps, diarrhea, and discomfort in the chest, call a physician and the National Poison Control Center hotline (1-800-222-1222).

Your call will be routed to your State Poison Control Center.

Anyone with a pesticide exposure poisoning emergency can call the toll-free telephone number for help. Personnel at the Center will give you first-aid information and direct you to local treatment centers if necessary.

For immediate medical attention call 911. Prompt action and treatment may save a life.



In Case of an Accident

- Remove the person from exposure.
- Get away from the treated or contaminated area immediately.
- Remove contaminated clothing.
- Wash with soap and clean water.
- Call a physician and the Poison Control Center (1-800-222-1222) or agency in your state.
- **Have the pesticide label with you! Follow the First Aid Precautionary Statements.**
- Be prepared to give the EPA registration number to the responding center/agency.